Hampshire Country Club Planned Residential Development Village of Mamaroneck, Westchester County, New York Draft Environmental Impact Statement

F Soil Resource Report






United States Department of Agriculture

NRCS

Natural Resources Conservation Service A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Westchester County, New York

Hampshire Country Club



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (http:// offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	
Soil Map	
Soil Map	
Legend	
Map Unit Legend	
Map Unit Descriptions	
Westchester County, New York	
CrC—Charlton-Chatfield complex, rolling, very rocky	
CtC—Chatfield-Hollis-Rock outcrop complex, rolling	
Uc—Udorthents, wet substratum	
Uf—Urban land	17
UIC—Urban land-Charlton-Chatfield complex, rolling, very rocky	18
W—Water	
References	21

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil scientists classified and named the soils in the survey area, they compared the

individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soillandscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



	MAP L	EGEND		MAP INFORMATION
Area of Intere	. ,	333	Spoil Area	The soil surveys that comprise your AOI were mapped at 1:12,000.
	rea of Interest (AOI)	٥	Stony Spot	Please rely on the bar scale on each map sheet for map
Soils	oil Map Unit Polygons	0	Very Stony Spot	measurements.
	oil Map Unit Lines	8	Wet Spot	Source of Map: Natural Resources Conservation Service
	oil Map Unit Points	\triangle	Other	Web Soil Survey URL: http://websoilsurvey.nrcs.usda.gov
Special Poi	·	, * *:	Special Line Features	Coordinate System: Web Mercator (EPSG:3857)
() BI	lowout	Water Fea		Maps from the Web Soil Survey are based on the Web Mercator
B	orrow Pit	~~ T	Streams and Canals	projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the
ж с	lay Spot	Transport	Rails	Albers equal-area conic projection, should be used if more accurate
♦ C	losed Depression	~	Interstate Highways	calculations of distance or area are required.
💥 G	iravel Pit	~	US Routes	This product is generated from the USDA-NRCS certified data as of
G	iravelly Spot	~	Major Roads	the version date(s) listed below.
🙆 La	andfill	~	Local Roads	Soil Survey Area: Westchester County, New York
۸. La	ava Flow	Backgrou	nd	Survey Area Data: Version 11, Sep 25, 2015
M ملله	larsh or swamp	and the second s	Aerial Photography	Soil map units are labeled (as space allows) for map scales 1:50,000
🙊 M	line or Quarry			or larger.
© M	liscellaneous Water			Date(s) aerial images were photographed: Jul 21, 2014—Aug 27,
O P	erennial Water			2014
v R	ock Outcrop			The orthophoto or other base map on which the soil lines were
+ S	aline Spot			compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting
s Si	andy Spot			of map unit boundaries may be evident.
i Se	everely Eroded Spot			
👌 Si	inkhole			
si 🔉	lide or Slip			
<i>j</i> ø So	odic Spot			

Westchester County, New York (NY119)					
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI		
CrC	Charlton-Chatfield complex, rolling, very rocky	7.7	7.2%		
CtC	Chatfield-Hollis-Rock outcrop complex, rolling	24.1	22.5%		
Uc	Udorthents, wet substratum	62.6	58.4%		
Uf	Urban land	0.0	0.0%		
UIC	Urban land-Charlton-Chatfield complex, rolling, very rocky	11.9	11.1%		
W	Water	0.9	0.8%		
Totals for Area of Interest	·	107.2	100.0%		

Map Unit Legend

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic classes rarely, if ever, can be mapped without including areas of other taxonomic classes for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An association is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Westchester County, New York

CrC—Charlton-Chatfield complex, rolling, very rocky

Map Unit Setting

National map unit symbol: bd8f Elevation: 100 to 1,000 feet Mean annual precipitation: 46 to 50 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 115 to 215 days Farmland classification: Not prime farmland

Map Unit Composition

Charlton and similar soils: 50 percent Chatfield and similar soils: 30 percent Minor components: 20 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton

Setting

Landform: Ridges, hills, till plains Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Acid loamy till derived mainly from schist, gneiss, or granite

Typical profile

H1 - 0 to 8 inches: loam H2 - 8 to 24 inches: sandy loam H3 - 24 to 60 inches: sandy loam

Properties and qualities

Slope: 2 to 15 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 7.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6s Hydrologic Soil Group: B

Description of Chatfield

Setting

Landform: Ridges, hills Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Loamy till derived mainly from granite, gneiss, or schist

Typical profile

H1 - 0 to 7 inches: loam H2 - 7 to 24 inches: flaggy silt loam H3 - 24 to 28 inches: unweathered bedrock

Properties and qualities

Slope: 2 to 15 percent
Depth to restrictive feature: 20 to 40 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Low to high (0.01 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 1 percent
Available water storage in profile: Low (about 3.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6s Hydrologic Soil Group: B

Minor Components

Hollis

Percent of map unit: 5 percent

Rock outcrop

Percent of map unit: 5 percent

Sutton

Percent of map unit: 4 percent

Sun

Percent of map unit: 2 percent Landform: Depressions

Leicester

Percent of map unit: 2 percent

Palms

Percent of map unit: 1 percent Landform: Marshes, swamps

Carlisle

Percent of map unit: 1 percent Landform: Marshes, swamps

CtC—Chatfield-Hollis-Rock outcrop complex, rolling

Map Unit Setting

National map unit symbol: bd8h Elevation: 100 to 1,000 feet Mean annual precipitation: 46 to 50 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 115 to 215 days Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 30 percent Chatfield and similar soils: 30 percent Rock outcrop: 20 percent Minor components: 20 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Chatfield

Setting

Landform: Ridges, hills Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Loamy till derived mainly from granite, gneiss, or schist

Typical profile

H1 - 0 to 7 inches: loam H2 - 7 to 24 inches: flaggy silt loam H3 - 24 to 28 inches: unweathered bedrock

Properties and qualities

Slope: 3 to 15 percent
Depth to restrictive feature: 20 to 40 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Low to high (0.01 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 1 percent
Available water storage in profile: Low (about 3.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6s Hydrologic Soil Group: B

Description of Hollis

Setting

Landform: Ridges, hills

Landform position (two-dimensional): Shoulder

Landform position (three-dimensional): Crest

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: A thin mantle of loamy till derived mainly from schist, granite, and gneiss

Typical profile

H1 - 0 to 1 inches: fine sandy loam

H2 - 1 to 16 inches: fine sandy loam

H3 - 16 to 20 inches: unweathered bedrock

Properties and qualities

Slope: 3 to 15 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Natural drainage class: Somewhat excessively drained
Capacity of the most limiting layer to transmit water (Ksat): Very low (0.00 to 0.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6s Hydrologic Soil Group: D

Description of Rock Outcrop

Properties and qualities

Slope: 3 to 15 percent
Depth to restrictive feature: 0 inches to lithic bedrock
Capacity of the most limiting layer to transmit water (Ksat): Low to very high (0.01 to 19.98 in/hr)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6s

Minor Components

Charlton

Percent of map unit: 8 percent

Sutton

Percent of map unit: 5 percent

Leicester

Percent of map unit: 2 percent

Sun

Percent of map unit: 2 percent Landform: Depressions

Unnamed soils, very shallow

Percent of map unit: 2 percent

Palms

Percent of map unit: 1 percent *Landform:* Marshes, swamps

Uc—Udorthents, wet substratum

Map Unit Setting

National map unit symbol: bd7g Elevation: 50 to 2,400 feet Mean annual precipitation: 46 to 50 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 115 to 215 days Farmland classification: Not prime farmland

Map Unit Composition

Udorthents, wet substratum, and similar soils: 80 percent *Minor components:* 20 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Udorthents, Wet Substratum

Typical profile

H1 - 0 to 4 inches: gravelly loam *H2 - 4 to 72 inches:* very gravelly loam

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: 40 to 60 inches to lithic bedrock
Natural drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high (0.06 to 5.95 in/hr)
Depth to water table: About 6 to 24 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 15 percent
Available water storage in profile: Low (about 4.6 inches)

Minor Components

Udorthents

Percent of map unit: 5 percent

Urban land

Percent of map unit: 5 percent

Fredon

Percent of map unit: 2 percent Landform: Depressions

Paxton

Percent of map unit: 2 percent

lpswich

Percent of map unit: 2 percent Landform: Tidal marshes

Raynham

Percent of map unit: 2 percent

Hinckley

Percent of map unit: 2 percent

Uf—Urban land

Map Unit Setting

National map unit symbol: bd7j Elevation: 50 to 2,400 feet Mean annual precipitation: 46 to 50 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 115 to 215 days Farmland classification: Not prime farmland

Map Unit Composition

Urban land: 85 percent *Minor components:* 15 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Minor Components

Udorthents

Percent of map unit: 5 percent

Riverhead

Percent of map unit: 2 percent

Udorthents, wet substratum

Percent of map unit: 2 percent

Unadilla

Percent of map unit: 2 percent

Chatfield

Percent of map unit: 2 percent

Sutton

Percent of map unit: 2 percent

UIC—Urban land-Charlton-Chatfield complex, rolling, very rocky

Map Unit Setting

National map unit symbol: bd7n Elevation: 100 to 1,000 feet Mean annual precipitation: 46 to 50 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 115 to 215 days Farmland classification: Not prime farmland

Map Unit Composition

Urban land: 40 percent Charlton and similar soils: 20 percent Chatfield and similar soils: 15 percent Minor components: 25 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton

Setting

Landform: Ridges, hills, till plains Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Acid loamy till derived mainly from schist, gneiss, or granite

Typical profile

H1 - 0 to 8 inches: loam H2 - 8 to 24 inches: sandy loam H3 - 24 to 60 inches: sandy loam

Properties and qualities

Slope: 2 to 15 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 7.5 inches)

Description of Chatfield

Setting

Landform: Ridges, hills Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Loamy till derived mainly from granite, gneiss, or schist

Typical profile

H1 - 0 to 7 inches: loam H2 - 7 to 24 inches: flaggy silt loam H3 - 24 to 28 inches: unweathered bedrock

Properties and qualities

Slope: 2 to 15 percent
Depth to restrictive feature: 20 to 40 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Low to high (0.01 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 1 percent
Available water storage in profile: Low (about 3.2 inches)

Minor Components

Leicester

Percent of map unit: 5 percent Landform: Depressions

Sutton

Percent of map unit: 5 percent

Udorthents

Percent of map unit: 5 percent

Rock outcrop

Percent of map unit: 5 percent

Hollis

Percent of map unit: 2 percent

Sun

Percent of map unit: 2 percent Landform: Depressions

Palms

Percent of map unit: 1 percent Landform: Marshes, swamps

W—Water

Map Unit Setting

National map unit symbol: bd7z Mean annual precipitation: 46 to 50 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 115 to 215 days Farmland classification: Not prime farmland

Map Unit Composition

Water: 100 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/soils/?cid=nrcs142p2 054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.


United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2_053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2 054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf



USDA Natural Resources Conservation Service

	ND	MAP INFORMATION		
rest (AOI) Backg	ground	The soil surveys that comprise your AOI were mapped at 1:12,000.		
Area of Interest (AOI)	Aerial Photography	Please rely on the bar scale on each map sheet for map measurements.		
Ig Polygons Very limited Somewhat limited		Source of Map: Natural Resources Conservation Service Web Soil Survey URL: http://websoilsurvey.nrcs.usda.gov Coordinate System: Web Mercator (EPSG:3857)		
Not limited Not rated or not available Ig Lines		Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.		
Very limited Somewhat limited		This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.		
Not limited Not rated or not available		Soil Survey Area: Westchester County, New York Survey Area Data: Version 11, Sep 25, 2015		
ng Points Very limited		Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.		
Somewhat limited		Date(s) aerial images were photographed: Jul 21, 2014—Aug 27, 2014		
Not limited Not rated or not available		The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background		
ires Streams and Canals		imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.		
tion Rails				
Interstate Highways				
US Routes				
Major Roads				
Local Roads				
	rest (AOI) Back Area of Interest (AOI) Back Very limited Somewhat limited Not rated or not available g Lines Very limited Somewhat limited Not rated or not available g Points Very limited Somewhat limited Not rated or not available g Points Very limited Somewhat limited Not rated or not available res Streams and Canals ion Rails Interstate Highways US Routes Major Roads	Area of Interest (AOI) Aerial Photography g Polygons Very limited Somewhat limited Not rated or not available g Lines Very limited Somewhat limited Not rated or not available g Points Very limited Somewhat limited Not rated or not available g Points Very limited Somewhat limited Not rated or not available g Points Very limited Somewhat limited Not rated or not available gress Streams and Canals ion Rails Interstate Highways US Routes Major Roads		

Dwellings With Basements

Map unit symbol	Map unit name	Rating	Component name (percent)	Rating reasons (numeric values)	Acres in AOI	Percent of AOI
CrC	Charlton- Chatfield complex, rolling, very rocky	Somewhat limited	Charlton (50%)	Slope (0.04)	7.7	7.2%
CtC Chatfield-Hollis- Rock outcrop	Very limited	Chatfield (30%)	Depth to hard bedrock (1.00)	24.1	22.5%	
	complex, rolling			Slope (0.04)		
			Hollis (30%)	Depth to hard bedrock (1.00)		
				Slope (0.04)		
Uc Udorthents, wet substratum	Very limited	Udorthents, wet substratum (80%)	Depth to saturated zone (1.00)	62.6	58.4%	
			Depth to hard bedrock (0.42)			
Uf	Urban land	nd Not rated	Urban land (85%)		0.0	0.0%
			Unadilla (2%)			
			Chatfield (2%)			
			Sutton (2%)			
			Riverhead (2%)			
UIC Urban land- Charlton- Chatfield complex, rolling, very rocky		Not rated	Urban land (40%)		11.9	11.1%
		Leicester (5%)				
		ng, very	Sutton (5%)			
			Udorthents (5%)			
			Rock outcrop (5%)			
			Hollis (2%)			
			Sun (2%)			
		Palms (1%)				
W	Water	Not rated	Water (100%)		0.9	0.8%
Totals for Area	of Interest				107.2	100.0%

Dwellings With Basements— Summary by Rating Value			
Rating Acres in AOI Percent of AO			
Very limited	86.7	80.9%	
Somewhat limited	7.7	7.2%	

Dwellings With Basements— Summary by Rating Value			
Rating	Acres in AOI	Percent of AOI	
Null or Not Rated	12.8	12.0%	
Totals for Area of Interest	107.2	100.0%	

Description

Dwellings are single-family houses of three stories or less. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet.

The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification of the soil. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. "Not limited" indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. "Somewhat limited" indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. "Very limited" indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The map unit components listed for each map unit in the accompanying Summary by Map Unit table in Web Soil Survey or the Aggregation Report in Soil Data Viewer are determined by the aggregation method chosen. An aggregated rating class is shown for each map unit. The components listed for each map unit are only those that have the same rating class as listed for the map unit. The percent composition of each component in a particular map unit is presented to help the user better understand the percentage of each map unit that has the rating presented.

Other components with different ratings may be present in each map unit. The ratings for all components, regardless of the map unit aggregated rating, can be viewed by generating the equivalent report from the Soil Reports tab in Web Soil Survey or from the Soil Data Mart site. Onsite investigation may be needed to validate these interpretations and to confirm the identity of the soil on a given site.

Rating Options

Aggregation Method: Dominant Condition

Component Percent Cutoff: None Specified

Tie-break Rule: Higher